



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, C07K 14/705, A61K 38/17, G01N 33/68	A1	(11) International Publication Number: WO 00/68377 (43) International Publication Date: 16 November 2000 (16.11.00)
(21) International Application Number: PCT/EP00/03683 (22) International Filing Date: 25 April 2000 (25.04.00) (30) Priority Data: 99109094.5 7 May 1999 (07.05.99) EP (71) Applicant (for all designated States except US): MERCK PATENT GMBH [DE/DE]; Frankfurter Str. 250, D-64293 Darmstadt (DE). (72) Inventor; and (75) Inventor/Applicant (for US only): CLEMETSON, Kenneth, J. [GB/CH]; Fluracker 29, CH-3065 Bern (CH). (74) Common Representative: MERCK PATENT GMBH; D-64271 Darmstadt (DE).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: RECOMBINANT PLATELET COLLAGEN RECEPTOR GLYCOPROTEIN VI AND ITS PHARMACEUTICAL USE (57) Abstract The invention relates to Glycoprotein VI (GPVI), its isolation, purification, and methods for recombinant production. Especially, the invention relates to the use of GPVI, preferably recombinant GPVI, in the treatment of disorders and pathological events correlated directly or indirectly to blood coagulation disorders such as thrombotic and cardiovascular diseases. The extracellular recombinant protein can also be used for establishing screening assays to find potential inhibitors of the membrane bound GPVI in order to inhibit binding of thrombocytes and platelets, respectively, to collagen. Changes in GPIV can be used to monitor platelet age and exposure to thrombotic and cardiovascular diseases.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

RECOMBINANT PLATELET COLLAGEN RECEPTOR GLYCOPROTEIN VI AND ITS PHARMACEUTICAL USE

- 5 The invention relates to Glycoprotein VI (GPVI), its isolation, purification, and methods for recombinant production. Especially, the invention relates to the use of GPVI, preferably recombinant GPVI, in the treatment of disorders and pathological events correlated directly or indirectly to blood coagulation disorders such as thrombotic and cardiovascular diseases. The extracellular recombinant
- 10 protein can also be used for establishing screening assays to find potential inhibitors of the membrane bound GPVI in order to inhibit interaction of platelets and collagen. GPVI on the platelet surface is modified during the platelet lifetime in vivo and can therefore be used as a marker of the platelet age profile.
- 15 Glycoprotein VI is a 62/65 kDa (non-reduced/reduced respectively) platelet membrane glycoprotein which forms a complex together with the Fc γ common subunit. The GPVI subunit contains the collagen binding site and the Fc γ subunit is responsible for signalling. The complex forms one of the major collagen receptors on the platelet surface, critical for platelet activation in response to
- 20 collagen. The recognition sequence on collagen consists of (GlyProHyp)_n sequences. Patients are known from Japan who have a genetic deficiency of GPVI. They have mild bleeding problems and their platelets respond only weakly to collagen, presumably via other receptors. A great deal has been learned about the signalling cascades originating at GPVI which strongly resemble those from
- 25 immune receptors including T-cell receptors, B-cell receptors and natural killer cell receptors. These cascades involve src family tyrosine kinases such as Fyn and Lyn as well as p72^{SYK} and many other tyrosine kinases and phosphatases and adaptor proteins such as LAT. A main target of these cascades is activation of phospholipase C γ 2 which splits phospholipids to give the second messengers
- 30 diacylglycerol and IP₃. GPVI is thought to be involved in activation of the platelet integrin α 2 β 1 which has a major role in platelet adhesion to damaged vessel wall. Mice with the Fc γ subunit "knocked-out" have platelets which still show responses to collagen implying that the resting state of α 2 β 1 may also be regulated by the GPVI/Fc γ complex.

The platelet collagen receptor GPVI is closely related to the natural killer activatory receptors of the p58KAR family as well as to Fc α R.

The adhesion and activation of resting, circulating platelets at a site of vascular injury is the first step in a process leading to the formation of a thrombus which is converted into a haemostatic plug. Collagen is one of the major components of the vessel wall responsible for platelet activation. Many types of collagen exist and seven of these are found in the subendothelial layers. Several different receptors for collagen have been identified on platelets but the major ones are now considered to be the integrin $\alpha_2\beta_1$ and the non-integrin GPVI. Although $\alpha_2\beta_1$ is well characterised and both subunits were cloned and sequenced several years ago, the structure of GPVI has remained elusive although several features have been identified. It was determined about twenty years ago that GPVI is a major platelet glycoprotein with a molecular mass in the 60-65 kDa range and an acid pI. Its role as a putative collagen receptor was established following the identification of a patient in Japan with a mild bleeding disorder whose platelets showed a specific defect of response to collagen and lacked this receptor. This patient had also developed autoantibodies to the deficient receptor and these were used to characterise the molecule further. More recently it was established that GPVI is associated non-covalently with the common Fc γ subunit which acts as the signalling part of the complex. It was also demonstrated that the recognition sequence on collagen for GPVI is a repeated Gly-Pro-Hyp triplet within the collagen triple helical structure and that synthetic peptides based on this structure could be used as specific GPVI directed agonists. The GPVI/ Fc γ complex was shown to signal to the platelet interior by an immune receptor-like mechanism, involving activation of p72^{SYK} and leading by a cascade of kinase/phosphatase/adaptor protein interactions to activation of PLC γ 2 and hence to release of granules and platelet aggregation. A further step in characterisation of this molecule was the demonstration that the snake C-type lectin, convulxin, from the Tropical Rattlesnake, *Crotalus durissus terrificus* was able to activate platelets by clustering GPVI through a multimeric interaction. Convulxin was shown to bind specifically to GPVI providing a method for purification of this receptor in conjunction with established approaches.

Thus, it is clear from the prior art that GPVI seems to be a very interesting compound in many therapeutical fields above all concerning with applications which are related, directly or indirectly, to blood coagulation events which depend on collagen – platelet interaction. It was, therefore, the goal of the present invention is to provide GPVI in a recombinant form and to show its efficiency as direct therapeutical target or as tool for screening of short compounds, especially chemically synthesized or synthesizable compounds having the capability to inhibit or block the natural platelet-collagen interaction.

The invention relates also to portions or fragments of the GPVI protein which have maintained their biological activity which is the binding to collagen.

The invention was successful in purifying adequate amounts of GPVI for preliminary characterisation and for peptide sequencing. The sequences were used to design primers for PCR to identify a positive sequence in a DNA library. This DNA sequence was then used as a probe to isolate an almost complete cDNA sequence from the library and missing 5'-sequence was obtained using a RACE method from a platelet cDNA library.

The invention was also successful in showing the use of recombinant GPVI as therapeutically applicable compound which is capable, when administered in a patient with e.g. damaged blood vessels, to bind to collagen, thus preventing platelets bearing membrane-bound GPVI from binding to said collagen.

The recombinant soluble extracellular domain of GPVI contains the collagen binding site and can prevent platelet activation by collagen. It could therefore be applicable to treatment of disease conditions involving increased platelet activation with collagen, such as atherosclerotic plaque rupture, in diseases such as unstable angina or, during surgical treatment such as Percutaneous Transluminal Coronary Angioplasty (PTCA), where arteries are reopened by inflation of a balloon catheter causing considerable damage to the vessel wall and much platelet activation and often resulting in reclosure of the vessel later. The advantage of recombinant GPVI fragments compared to present treatment methods is that they act at an earlier stage by preventing or reducing platelet activation rather than by suppressing events after platelet activation, such as

aggregation by GPIIb-IIIa antagonists. Thus, smaller amounts of platelet granule contents are released including growth factors and chemokines which are involved not only in wound repair but in the remodelling of the vessel wall by smooth muscle migration and in attraction of phagocytic cells such as monocytes
5 known to contribute to atherosclerosis. Fab fragment of humanised mouse monoclonal antibodies against GPVI could be used with similar effect to block GPVI on the platelet surface with similar applications as above.

Recombinant GPVI according to this invention can also be used in a binding
10 assay to collagen to screen for small molecules (in combinatorial libraries for example) capable of inhibiting this interaction and which can be used to develop therapeutic compounds which are inhibitors of the collagen-platelet interaction . By suitable derivatisation these compounds are made orally available. Again the main objective is to prepare compounds reducing GPVI-collagen interactions and
15 hence platelet activation in situations where platelets come into contact with collagen. The screening technology as such used in this invention is well established in the prior art. By such screening assays the invention enables finding and developing new targets which can inhibit the natural membrane-bound GPVI on the platelet surface as a collagen antagonist. Such targets which
20 may be small chemical molecules may then be the basis for further inventions.

Another major application of GPVI and reagents that recognize specific domains of GPVI is as markers of platelet age and functionality. Young platelets are generally thought to be more active and functional than older ones. Young
25 platelets bind to and are activated by the snake venom C-type lectin convulxin, which is specific for GPVI, and as they age both the binding and degree of activation decrease. This can be due to either proteolytic or conformational changes in GPVI or its association with Fcγ due to platelet activation or damage in the circulation. This can be a useful parameter to measure the age and
30 function profile of platelets in patients as well as in normal persons during medical controls. The platelet age profile changes in many diseases affecting the bone marrow or the immune system and could be an important diagnostic criterion if better methods for its determination were available. For example, patients with

diseases involving increased platelet turnover will show more young platelets whereas patients on chemotherapy or radiation treatment will show a steadily aging population. Thus, such an age profile can be used for a precise monitoring of treatment. In a normal healthy population very little is known about the age profile distribution and its role as a predictor of changes in health. It is not known whether the changes in GPVI are due to the partial involvement of platelets in haemostatic events and whether changes might be more pronounced in patients with extensive cardiovascular disease. At present thiazole orange is used to detect young reticulated platelets containing mRNA. This mRNA soon decays, restricting the method to only the youngest platelets. Reagents which could be used in such an assay would include GPVI-specific snake venom proteins such as convulxin, or monoclonal or polyclonal antibodies recognising the N-terminal region of GPVI or monoclonal antibodies recognising new sites or conformations exposed by proteolysis of the N-terminal domain or specific conformations present either in the intact molecule and not in the aged one or vice versa or small chemical entities selected to recognise specifically intact GPVI or its modified form. These reagents would be labelled with a fluorescent marker, or together with a fluorescent labelled second antibody or affinity reagent and used in flow cytometry to measure the platelet binding profile. At a later stage alternative, less labour intensive measuring techniques based on automated measuring of platelet profiles could be adopted. Using cell sorting methods with flow cytometry or magnetic beads it should be possible to isolate young and old platelets to examine the factors involved in removal of old platelets from the circulation. Reagents recognizing specific forms of GPVI would be a key to such studies.

Therefore, it is an object of the present invention to provide a DNA coding for Glycoprotein VI or biological active fragments thereof, especially the sequence of Fig. 2.

It is a further object of this invention to provide a DNA coding for Glycoprotein VI comprising the amino acid sequences of Fig. 1a and 1b.

It is another object of this invention to provide a pharmaceutical composition comprising recombinant GPVI together with a pharmaceutically acceptable diluent, carrier or excipient, and its use for the manufacture of a medicament in the therapeutical field of thrombotic and cardiovascular events and disorders related to platelet-collagen interactions

Another object of the invention is the use of recombinant GPVI in a screening tool for detecting specific inhibitors of platelet-collagen interactions.

Another object of the invention is the use of GPVI as a marker for platelet age and exposure to cardiovascular diseases.

10

Possible medical indications and applications, respectively, are, for example, unstable angina pectoris, PTCA, use of stents in this field, operations on coronary vessels, general operations on blood vessels, operations which may damage larger blood vessels such as hip joint operations. Moreover, all indications are included which relate to thromboembolic events caused by disorders of the interaction between the vessel wall and the coagulation system with a high risk of formation of thrombi and blocking of vessels.

15

As indicated above, the GPVI protein and fragments thereof according to the present invention are suitable as pharmaceutically effective compounds in pharmaceutical compositions and combinations.

20

The pharmaceutical formulations according to the invention optionally may comprise additional active ingredients like anti-coagulants such as hirudin or heparin or thrombolytic agents such as plasminogen activator or hementin or antagonists to other platelet receptors such as GPIIb-IIIa antagonists like abciximab or eptifibatide or ADP-receptor antagonists such as clopidogrel.

25

The novel protein, and its biological active fragments respectively, according to the invention may form pharmaceutically acceptable salts with any non-toxic, organic or inorganic acid. Inorganic acids are, for example, hydrochloric, hydrobromic, sulphuric or phosphoric acid and acid metal salts such as sodium monohydrogen orthophosphate and potassium hydrogen sulfate. Examples for

30

organic acids are the mono, di and tri carboxylic acids such as acetic, glycolic, lactic, pyruvic, malonic, succinic, glutaric, fumaric, malic, tartaric, citric, ascorbic, maleic, hydroxymaleic, benzoic, hydroxybenzoic, phenylacetic, cinnamic, salicylic and sulfonic acids such as methane sulfonic acid. Salts of the carboxy terminal amino acid moiety include the non-toxic carboxylic acid salts formed with any suitable inorganic or organic bases. These salts include, for example, alkali metals such as sodium and potassium, alkaline earth metals such as calcium and magnesium, light metals of Group IIIA including aluminium, and organic primary, secondary and tertiary amines such as trialkylamines, including triethylamine, procaine, dibenzylamine, 1-ethenamine, N,N'-dibenzylethylene-diamine, dihydroabietylamine and N-alkylpiperidine.

As used herein, the term "pharmaceutically acceptable carrier" means an inert, non toxic solid or liquid filler, diluent or encapsulating material, not reacting adversely with the active compound or with the patient. Suitable, preferably liquid carriers are well known in the art such as sterile water, saline, aqueous dextrose, sugar solutions, ethanol, glycols and oils, including those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil and mineral oil.

The formulations according to the invention may be administered as unit doses containing conventional non-toxic pharmaceutically acceptable carriers, diluents, adjuvants and vehicles which are typical for parenteral administration.

The term "parenteral" includes herein subcutaneous, intravenous, intra-articular and intratracheal injection and infusion techniques. Also other administrations such as oral administration and topical application are suitable. Parenteral compositions and combinations are most preferably administered intravenously either in a bolus form or as a constant fusion according to known procedures.

Tablets and capsules for oral administration contain conventional excipients such as binding agents, fillers, diluents, tableting agents, lubricants, disintegrants, and wetting agents. The tablets may be coated according to methods well known in the art.

Oral liquid preparations may be in the form of aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for reconstitution with water or another suitable vehicle before use. Such liquid preparations may contain conventional additives like suspending agents, emulsifying agents, non-aqueous vehicles and preservatives. Topical applications may be in the form of aqueous or oily suspensions, solutions, emulsions, jellies or preferably emulsion ointments.

Unit doses according to the invention may contain daily required amounts of the protein according to the invention, or sub-multiples thereof to make up the desired dose. The optimum therapeutically acceptable dosage and dose rate for a given patient (mammals, including humans) depends on a variety of factors, such as the activity of the specific active material employed, the age, body weight, general health, sex, diet, time and route of administration, rate of clearance. the object of the treatment, i.e., therapy or prophylaxis and the nature of the thrombotic disease to be treated, antiplatelet or anticoagulant activity.

Therefore, in compositions and combinations useful as anticoagulants in a treated patient (in vivo) a pharmaceutical effective daily dose of the peptides of this invention is between about 0.01 and 100 mg/kg body weight, preferably between 0.1 and 10 mg/kg body weight. According to the application form one single dose may contain between 0.5 and 10 mg of the collagen inhibitor To achieve an anticoagulant effect in extracorporeal blood a pharmaceutically effective amount of the inventive peptides is between 0.2 and 150 mg /l, preferably between 1mg and 20 mg/l extracorporeal blood.

Short Description of the Figures:

Fig. 1: Protein sequence of GPVI (one-letter-code)

1a: Leader sequence

1b: Mature protein

Open reading frame: 339 amino acids

Asterisk: Glycosylation site
Double underline: Transmembrane domain
Underline: Sequenced peptides

- 5 Fig. 2: GPVI nucleotide sequence covering open reading frame of 1017 bp plus 3' and 5' regions total 1249 bp

Detailed description of the invention

Two sequences of 7 amino acids showing the least degeneracy in the genetic
10 code were chosen for the synthesis of DNA primers in order to amplify part of the
of GPVI cDNA by PCR. As the location of both peptides in the protein were totally
unknown, for each of them, two degenerate primers, one sense and one
antisense were prepared. These primers were used to amplify a human bone-
marrow library. The combination of the sense 5' TYA THC CNG CNA TGA ARMG
15 3' primer coding for the sequence PAMKRSL with the antisense 5' TTR TAN ARN
GCR AAY TGR TC 3' one corresponding to DQFALYK amplified a DNA fragment
of 221 bp. In addition to the selected peptides, the amplified DNA coded for the
LysC/AspN peptide DQLELVATGVFAKPSLSAQPGPAVSS, clearly linking the
sequence to the cDNA for GPVI.

20

Screening 600.000 pfu from a bone marrow library with this 221 bp DNA fragment
produced 4 positive pfu. Three had inserts of 1350 bp whether cut by the
restriction enzymes Sal I or EcoR I and belonged to the IgG superfamily. The
fourth one had an 4.6 kb insert by Sal I digestion and gave two fragments of 2300
25 bp and 1300 bp respectively when treated by EcoRI. Its DNA encoded the
sequence for the 10 peptides derived from amino acid sequencing of GPVI but
stopped short of the amino terminal. No starting methionine or leader sequence
could be found but more than 2000 bp of previously sequenced non-reading
frame DNA terminating in an Alu sequence were present. The 5' end RACE
30 experiment was completed on platelet poly A RNA with primers located in a part
of the GPVI sequence which had been corroborated by that of the peptides. A
fragment of 348 bp including 278 bp on the sequence of the fourth clone and 70
bp new from bp 1987 corresponding to 14 amino acids including the first

methionine were found before falling back on the established GPVI sequence. Thus, a cDNA containing a total of 1249 bp, a 25 bp 5' sequence upstream of the start codon, an open reading frame of 1017 bp coding for a protein, including leader sequence, with 339 amino acids, and a 3' region of 207 bp including the stop codon could be sequenced.

A cDNA coding for platelet GPVI was cloned and sequenced from a human bone marrow cDNA library using RACE with platelet mRNA to supply missing 5' sequence. The open reading frame of 1017 bp encodes 339 amino acids and a untranslated 3' region. Hydrophobicity analysis of the amino acid sequence revealed the presence of two putative transmembrane domains, a putative 20 amino acid signal sequence, and a 19 amino acid domain between residues 247 and 265 of the mature protein. The sequence and its amino acid translation are shown in Fig. 2 and Fig. 1. A comparison with the amino acid sequence of the most similar molecules found in a search of GenBank reveals clearly that it belongs to the immunoglobulin superfamily and the extracellular domain contains two Ig C2-domain loops formed by two disulphide bridges. It is a membrane crossing protein class one molecule with the N-terminus at the exterior and traverses the membrane once. The most closely related molecules belong to the natural killer receptor class which contains both inhibitory and activatory types. GPVI clearly belongs to the activatory subclass not only through its function but also because unlike the inhibitory class it does not contain ITIM sequences in its cytoplasmic domain. Neither does it contain any tyrosine residues which might be involved in phosphorylation. There are some threonine and serine residues in this domain but they do not match any criteria for kinase consensus sequences. Like the activatory class of NK receptors, GPVI contains an arginine residue as the third amino acid of the membrane crossing domain which is involved in the complex formation with the Fc γ subunit. The cytoplasmic domain contains 51 amino acids, showing only a minor similarity (in the region just below the membrane) to the cytoplasmic domains of other members of this family. This suggests that this domain in GPVI may associate with different types of cytoplasmic molecule than the other family members. GPVI contains only a single putative N-glycosylation site at Asn69. The domain just above the membrane

- after the beta sheets of the Ig domains finish, however, is rich in threonine and serine residues which could provide O-glycosylation sites such as are found in GPIb α and GPV. The main function of this O-glycosylation seems to be to present the receptor structures well-extended from the platelet surface to facilitate the interactions with their bulky ligands. Since GPVI was earlier established as a sialoglycoprotein, the difference in molecular mass between the theoretical amino acid mass (37 kDa) and the mass determined by gel electrophoresis (65 kDa reduced) must be due to this glycosylation.
- 10 The structure of natural killer receptors of the two domain type has been established by X-ray crystallographic studies and the two Ig-domains were shown to form an acute angle with the receptor site for the peptide-carrying HLA antigens lying on the outside of the elbow. A direct comparison of the structure of the HLA peptide binding site with that of collagen immediately suggests why
- 15 these receptors have a common origin because the multiple alpha-helical structures of the HLA binding site and the peptide it contains strongly resemble the triple helical structure of collagen. The natural killer receptors are postulated to work by a dimerisation mechanism with two receptors recognising two separate HLA sites on the cell which the natural killer cell is investigating.
- 20 Possibly this dimerisation is part of the activation or deactivation mechanism, depending on the class of receptor. In the case of GPVI there may as well be the possibility for two GPVI molecules to associate with one Fc γ , since each monomer of the Fc γ dimer has a recognition sequence. However, the stoichiometry is not yet known, and based upon the structure of collagens,
- 25 collagen-like peptides that act via GPVI and convulxin, it seems likely that the strength of the signal is related to the number of GPVI/Fc γ complexes that are clustered together. Other platelet receptors belonging to this Ig family include ICAM-2 (CD102) and PECAM (CD31).
- 30 All microorganisms, cell lines, expression systems, expression hosts, plasmids, promoters, resistance markers, replication origins, restriction sites or other fragments or parts of vectors which are mentioned in the description not directly in connection with the claimed invention are commercially or otherwise generally

available. Provided that no other hints are given, they are used only as examples and are not essential with respect to the invention, and can be replaced by other suitable tools and biological materials, respectively.

- 5 The techniques which are essential according to the invention are described in detail below and above. Other techniques which are not described in detail correspond to known standard methods which are well known to a person skilled in the art, or are described more in detail in the cited references and patent applications and in the standard literature (e.g. Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd Edition, Cold Spring Harbor; Harlow, Lane, 10 1988, Antibodies: A Laboratory Manual, Cold Spring Harbor).

EXAMPLES

- 15 Example 1: *Materials*—Protein A-Sepharose, peroxidase-conjugated goat anti-mouse and anti-rabbit antibodies, bovine serum albumin, *Crotalus durissus terrificus* venom, wheat germ agglutinin (WGA), N-hydroxysuccinimidyl chloroformate-activated cross-linked 4% beaded agarose and Triton X-114 were from Sigma Chemical Co. (St Louis, MO), Octanoyl-N-methyl-glucamide (ONMG) 20 and nonanoyl-N-methyl-glucamide (NNMG) were from Oxyl Chemie (Bobingen, Germany).

- Example 2: *GPVI isolation from platelets* – Membrane glycoproteins were isolated from platelets as previously described. Briefly, platelets (from 40 buffy 25 coats) were washed and lysed in 2% Triton X-114 in the presence of protease inhibitors. The Triton X-114 and aqueous phases were separated and the detergent phase was loaded on a column of wheat-germ agglutinin coupled to Sepharose 4B. The platelet glycoproteins were eluted with 10 mM Tris HCl, pH 7.4, 30 mM NaCl, 0.2% octanoyl-N-methylglucamide (ONMG) and 2% N- 30 acetylglucosamine. After dialysis and concentration, the solution of glycoproteins was loaded on a column of convulxin bound to N-hydroxylsuccinamidyl-p-nitrophenyl chloroformate activated cross-linked 4% beaded agarose (1 mg/ml). The column was washed with 4 volumes of 10 mM Tris HCl, pH7.4, 30 mM NaCl,

0.2% nonanoyl-N-methylglucamide (NNMG), and then with 4 volumes of 10 mM Tris HCl, pH7.4, 30 mM NaCl and 2% NNMG. GPVI was eluted with 0.08% SDS in 10 mM Tris/HCl, pH 7.4. The solution was concentrated and loaded on a preparative gel of 8.5 % polyacrylamide using the Model 491 Prep Cell (BioRad, CA). The preparative electrophoresis was performed under non-reduced conditions following the manufacturer instructions. GPVI eluted as a single band at 65 kDa. The fractions were pooled, concentrated on Centricon-30 (Amicon, Beverly, MA) and resuspended in 10 mM Tris/HCl, pH7.4 and 0.1% ONMG.

10 Example 3: *Amino acid analysis of GPVI* - GPVI was digested with the endoproteinases LysC and AspN (Boehringer Mannheim, Germany). The 10 peptides generated were separated by reverse-phase HPLC and sequenced on an Applied Biosystem model 477A pulsed-liquid-phase protein sequencer with a model 120A on-line phenylthiohydantoin amino acid analyser.

15

Example 4: *Amplification of a 221 bp fragment coding for part of GPVI from a λ gt11 cDNA library* -

A sample (10^{10} pfu) (plaque forming units) from a human bone marrow library (Clonotech, Palo Alto, CA) was amplified using 2 combinations of 4 degenerate primers. The final primer concentrations were 2 μ M, the dNTP concentration was 200 μ M and 2 U/100 μ l reaction AmpliTaq Gold (Perkin Elmer, Rotkreuz, Switzerland) were used. The PCR conditions were 5 cycles at 37°C followed by 30 cycles at 44°C. The sense 19mer 5' TYATHCCNGCNATGAARMG 3' and the antisense 20mer 5' TTRTANARNGCRAAYTGRTC 3' amplified a 221 bp fragment which was subcloned in Bluescript KS⁺ (Stratagene, La Jolla, CA) and sequenced using the T7 Sequenase kit (Amersham, Switzerland).

Example 5: *Screening the λ gt11 cDNA library with the 221 bp GPVI probe*- The 221 bp fragment was cut from the plasmid, cleaned and labelled with α^{32} P-ATP (20MBq/50 μ l, Hartmann Analytik, Braunschweig, Germany) using the High Prime Labelling kit (Boehringer Mannheim, Switzerland). The human bone marrow library was screened following the manufacturer instructions. Positive phages were grown, their DNA isolated and subcloned in BlueScript using either EcoRI or

Sal I sites and sequenced. Sequencing was performed using the ABS system of RACE- Platelet poly A RNA was prepared as previously described (Power et al., Cytokine 7, 479-482, 1995). Reverse transcription (30 µl) was performed using 5 µg of poly A RNA with the primer 5'TGAATGAGACGGTCAGTTCAGC 3' (20 µM),
5 dNTP (1mM), RNAsin (40 U), 1X AMV buffer and 20 U AMV reverse transcriptase for 20 min at 45°C followed by 20 min at 52°C. The reaction mixture was treated with 2 µl 6N NaOH at 65° C for 30 min, neutralised with 2µl 6N acetic acid, and concentrated in a Centricon 30 (Amicon). An anchor was ligated to the first strand DNA following the protocol of Aptes and Siebert (BioTechniques 15:
10 890-893, 1993). Nested PCR was performed using a primer complementary to the anchor and the primer 5' TTGTACAGAGCAAATTGGTC 3' (35 cycles, 55°C) and followed by the primer 5' GACCAGAGGCTTCCGTTCTG 3' (30 cycles at 53°C). The highest band (350 bp) was separated by agarose electrophoresis from the lower ones, subcloned into BlueScript, and sequenced.

15

Example 6: *Preparation of anti-GPVI Fab and F(ab')₂*—Polyclonal antisera against human GPVI were generated in rabbits. IgG from rabbit anti-GPVI antiserum was purified as described. Digestion of IgG with immobilized papain (Pierce) to generate Fab fragments was performed according to the standard
20 protocol of the supplier. Fab fragments were separated from undigested IgG and Fc fragments using an immobilized Protein A (Sigma) column. The flowthrough was transferred to a dialysis tube, concentrated using solid polyethyleneglycol 20,000, thoroughly dialysed against 20 mM Hepes, 140 mM NaCl, 4 mM KCl, pH 7.4 and stored at 4°C until used. F(ab')₂ fragments were prepared by pepsin
25 digestion of IgG, 1:50 enzyme to substrate ratio (w/w), in 0.5 M acetate buffer, pH 4.0, at 37°C for 18 hr. The pH was corrected to 7.4 with diluted NaOH and the sample was dialysed against 20 mM phosphate, pH 7.4. F(ab')₂ fragments were separated from undigested IgG and Fc fragments using Protein A chromatography. The flow-through was transferred to dialysis tube, concentrated
30 using solid polyethyleneglycol 20 000, intensively dialysed against 20 mM Hepes, 140 mM NaCl, 4 mM KCl, pH 7.4 and stored in aliquots at -20°C. Washed platelets were lysed in Triton X-114 and phase separation was performed on the

soluble material before isolating the membrane glycoproteins associated with the Triton X-114 phase by affinity chromatography on wheat germ agglutinin-Sepharose 4B as described previously. As GPVI represents a very small fraction of the platelet membrane glycoprotein pool, we used the specificity of the snake
5 C-type lectin convulxin for isolation of this receptor. Affinity chromatography on convulxin coupled to Sepharose 4B yielded a 65 kDa protein as major product. However, uncharacterized material of both higher and lower Mr co-eluted with GPVI and could not be removed by extensive washing of the column. Preparative gel electrophoresis on 8.5 % polyacrylamide was added as a final step of
10 purification. Fractions containing GPVI were pooled and gave a single band on reanalysis. Purified GPVI was tested for its ability to block platelet aggregation by collagen. A slight inhibitory effect was observed when aliquots of GPVI solution were added to the platelet suspension. However, by preincubating GPVI with collagen before adding the mixture to the platelet suspension, aggregation could
15 be inhibited in a dose-dependant manner. These platelets still aggregated when fresh collagen was added. Under non-reducing conditions, the isolated protein has a Mr of 62 kDa with a shift toward a slightly higher Mr (65 kDa) under reducing conditions. As the amino terminus of GPVI was found to be blocked, the protein was digested with the enzymes LysC and LysC/AspN which produced 4
20 and 6 peptides, respectively, from which sequence was obtained. The peptides were separated by reverse phase HPLC on a C4 column and sequenced using the Edman method. The amino acid sequences of these peptides are underlined in the translated cDNA sequence (Fig. 1).

Patent claims

1. DNA coding for Glycoprotein VI or biologically active fragments thereof.
5
2. DNA according to claim 1 comprising partially or completely the sequence of Fig. 2.
3. DNA having the sequence of Fig. 2
10
4. DNA according to claims 1 – 3 coding for Glycoprotein VI comprising the amino acid sequence of Fig. 1a and 1b.
5. Recombinant human Glycoprotein VI as medicament comprising the amino
15 acid sequence of Fig. 1b.
6. Pharmaceutical composition comprising the protein of claim 5 and a pharmaceutically acceptable diluent, carrier or excipient therefor.
- 20 7. Pharmaceutical composition comprising additionally a pharmacologically active compound.
8. Use of recombinant GPVI in a screening tool for detecting specific inhibitors of platelet-collagen interactions.
25
9. Use of recombinant GPVI for the manufacture of a medicament in the therapeutic field of thrombotic and cardiovascular events and disorders related to platelet-collagen interactions.
- 30 10. Use of changes in GPVI as a marker for platelet age and platelet exposure to thrombotic and cardiovascular conditions or events.

Figure 1.(1a)

1 20

MSPSP~~TALFC~~ LGLCLGRVPA(1b)

1

QSGPLPKPSL QALPSSLVPL EKPVTLCQG PPGVDLYRLE KLSSRYQDQ (50)AVLFIPAMKR SLAGRYRCSY QN*GSLWSLPS DQLELVATGV FAKPSLSAQP (100)GPAVSSGGDV TLQCQTRYGF DQFALYKEGD PAPYKNPERW YRASFPIIITV (150)TAAHSGTYRC YSFSSRDPYL WSAPSDPLEL VVTGTSVTPS RLPTEPPSSV (200)AEFSEATAEL TVSFTNKVFT TETSR~~SITTS~~ PKESDSPAGP ARQYYTKGNL (250)VRICLGAVIL IILAGFLAED WHSRRKRLRH RGRAVQRPLP PLPPLPQTRK (300)SHGGQDGGRO DVHSRGLCS (319)

Figure 2:

GAGCTCAGGA CAGGGCTGAG GAACCATGTC TCCATCCCCG ACCGCCCTCT (50)
TCTGTCTTGG GCTGTGTCTG GGGCGTGTGC CAGCGCAGAG TGGACCGCTC (100)
CCCAAGCCCT CCCTCCAGGC TCTGCCCAGC TCCCTGGTGC CCCTGGAGAA
GCCAGTGACC CTCCGGTGCC AGGGACCTCC GGGCGTGGAC CTGTACCGCC (200)
TGGAGAAGCT GAGTTCCAGC AGGTACCAGG ATCAGGCAGT CCTCTTCATC
CCGGCCATGA AGAGAAGTCT GGCTGGACGC TACCGCTGCT CCTACCAGAA (300)
CGGAAGCCTC TGGTCCCTGC CCAGCGACCA GCTGGAGCTC GTTGCCACGG
GAGTTTTTGC CAAACCCTCG CTCTCAGCCC AGCCCGGCCC GCGGGTGTCTG (400)
TCAGGAGGGG ACGTAACCCT ACAGTGTGAG ACTCGGTATG GCTTTGACCA
ATTTGCTCTG TACAAGGAAG GGGACCCTGC GCCCTACAAG AATCCCGAGA (500)
GATGGTACCG GGCTAGTTTC CCCATCATCA CGGTGACCGC CGCCCACAGC
GGAACCTACC GATGCTACAG CTTCTCCAGC AGGGACCCAT ACCTGTGGTC (600)
GGCCCCCAGC GACCCCCTGG AGCTTGTGGT CACAGGAACC TCTGTGACCC
CCAGCCGGTT ACCAACAGAA CCACCTTCCT CGGTAGCAGA ATTCTCAGAA (700)
GCCACCGCTG AACTGACCGT CTCATTCACA AACAAAGTCT TCACAACTGA
GACTTCTAGG AGTATCACCA CCAGTCCAAA GGAGTCAGAC TCTCCAGCTG (800)
GTCCTGCCCCG CCAGTACTAC ACCAAGGGCA ACCTGGTCCG GATATGCCTC
GGGGCTGTGA TCCTAATAAT CCTGGCGGGG TTTCTGGCAG AGGACTGGCA (900)
CAGCCGGAGG AAGCGCCTGC GGCACAGGGG CAGGGCTGTG CAGAGGCCGC
TTCCGCCCCCT GCCGCCCCTC CCGCAGACCC GGAAATCACA CGGGGGTCAG (1000)
GATGGAGGCC GACAGGATGT TCACAGCCGC GGGTTATGTT CATGACCGCT
GAACCCAGG CACGGTCGTA TCCAAGGGAG GGATCATGGC ATGGGAGGCG (1100)
ACTCAAAGAC TGGCGTGTGT GGAGCGTGGA AGCAGGAGGG CAGAGGCTAC
AGCTGTGGAA ACGAGGCCAT GCTGCCTCCT CCTGGTGTTT CATCAGGGAG (1200)
CCGTTCCGGCC AGTGTCTGTC TGTCTGTCTG CCTCTCTGTC TGAGGGCAC (1249)

SEQUENCE LISTING

<110> Merck Patent GmbH

5 <120> Recombinant platelet collagen receptor glycoprotein VI
and its pharmaceutical use

<130> GlycoproteinVI-Merck

10 <140>
<141>

<160> 1

15 <170> PatentIn Ver. 2.1

<210> 1
<211> 1249
<212> DNA

20 <213> Homo sapiens

<220>
<221> CDS
<222> (26)..(85)

25 <223> Leader sequence

<220>
<221> CDS
<222> (86)..(1042)

30 <223> mature GPVI protein

<400> 1
gagctcagga cagggctgag gaacc atg tct cca tcc ccg acc gcc ctc ttc 52
Met Ser Pro Ser Pro Thr Ala Leu Phe

35 1 5

tgt ctt ggg ctg tgt ctg ggg cgt gtg cca gcg cag agt gga ccg ctc 100
Cys Leu Gly Leu Cys Leu Gly Arg Val Pro Ala Gln Ser Gly Pro Leu

40 10 15 20 25

ccc aag ccc tcc ctc cag gct ctg ccc agc tcc ctg gtg ccc ctg gag 148
Pro Lys Pro Ser Leu Gln Ala Leu Pro Ser Ser Leu Val Pro Leu Glu

30 35 40

45 aag cca gtg acc ctc cgg tgc cag gga cct ccg ggc gtg gac ctg tac 196
Lys Pro Val Thr Leu Arg Cys Gln Gly Pro Pro Gly Val Asp Leu Tyr

45 50 55

cgc ctg gag aag ctg agt tcc agc agg tac cag gat cag gca gtc ctc 244
50 Arg Leu Glu Lys Leu Ser Ser Ser Arg Tyr Gln Asp Gln Ala Val Leu

60 65 70

ttc atc ccg gcc atg aag aga agt ctg gct gga cgc tac cgc tgc tcc 292
Phe Ile Pro Ala Met Lys Arg Ser Leu Ala Gly Arg Tyr Arg Cys Ser

55 75 80 85

tac cag aac gga agc ctc tgg tcc ctg ccc agc gac cag ctg gag ctc 340
Tyr Gln Asn Gly Ser Leu Trp Ser Leu Pro Ser Asp Gln Leu Glu Leu

90 95 100 105

60

	gtt gcc acg gga gtt ttt gcc aaa ccc tcg ctc tca gcc cag ccc ggc	388
	Val Ala Thr Gly Val Phe Ala Lys Pro Ser Leu Ser Ala Gln Pro Gly	
	110 115 120	
5	ccg gcg gtg tcg tca gga ggg gac gta acc cta cag tgt cag act cgg	436
	Pro Ala Val Ser Ser Gly Gly Asp Val Thr Leu Gln Cys Gln Thr Arg	
	125 130 135	
10	tat ggc ttt gac caa ttt gct ctg tac aag gaa ggg gac cct gcg ccc	484
	Tyr Gly Phe Asp Gln Phe Ala Leu Tyr Lys Glu Gly Asp Pro Ala Pro	
	140 145 150	
	tac aag aat ccc gag aga tgg tac cgg gct agt ttc ccc atc atc acg	532
15	Tyr Lys Asn Pro Glu Arg Trp Tyr Arg Ala Ser Phe Pro Ile Ile Thr	
	155 160 165	
	gtg acc gcc gcc cac agc gga acc tac cga tgc tac agc ttc tcc agc	580
	Val Thr Ala Ala His Ser Gly Thr Tyr Arg Cys Tyr Ser Phe Ser Ser	
20	170 175 180 185	
	agg gac cca tac ctg tgg tcg gcc ccc agc gac ccc ctg gag ctt gtg	628
	Arg Asp Pro Tyr Leu Trp Ser Ala Pro Ser Asp Pro Leu Glu Leu Val	
	190 195 200	
25	gtc aca gga acc tct gtg acc ccc agc cgg tta cca aca gaa cca cct	676
	Val Thr Gly Thr Ser Val Thr Pro Ser Arg Leu Pro Thr Glu Pro Pro	
	205 210 215	
30	tcc tcg gta gca gaa ttc tca gaa gcc acc gct gaa ctg acc gtc tca	724
	Ser Ser Val Ala Glu Phe Ser Glu Ala Thr Ala Glu Leu Thr Val Ser	
	220 225 230	
	ttc aca aac aaa gtc ttc aca act gag act tct agg agt atc acc acc	772
35	Phe Thr Asn Lys Val Phe Thr Thr Glu Thr Ser Arg Ser Ile Thr Thr	
	235 240 245	
	agt cca aag gag tca gac tct cca gct ggt cct gcc cgc cag tac tac	820
	Ser Pro Lys Glu Ser Asp Ser Pro Ala Gly Pro Ala Arg Gln Tyr Tyr	
40	250 255 260 265	
	acc aag ggc aac ctg gtc cgg ata tgc ctc ggg gct gtg atc cta ata	868
	Thr Lys Gly Asn Leu Val Arg Ile Cys Leu Gly Ala Val Ile Leu Ile	
	270 275 280	
45	atc ctg gcg ggg ttt ctg gca gag gac tgg cac agc cgg agg aag cgc	916
	Ile Leu Ala Gly Phe Leu Ala Glu Asp Trp His Ser Arg Arg Lys Arg	
	285 290 295	
50	ctg cgg cac agg ggc agg gct gtg cag agg ccg ctt ccg ccc ctg ccg	964
	Leu Arg His Arg Gly Arg Ala Val Gln Arg Pro Leu Pro Pro Leu Pro	
	300 305 310	
	ccc ctc ccg cag acc cgg aaa tca cac ggg ggt cag gat gga ggc cga	1012
55	Pro Leu Pro Gln Thr Arg Lys Ser His Gly Gly Gln Asp Gly Gly Arg	
	315 320 325	
	cag gat gtt cac agc cgc ggg tta tgt tca tgaccgctga accccaggca	1062
	Gln Asp Val His Ser Arg Gly Leu Cys Ser	
60	330 335	

cggtcgtatc caagggaggg atcatggcat gggaggcgac tcaaagactg gcgtgtgtgg 1122
 agcgtggaag caggagggca gaggctacag ctgtggaaac gaggccatgc tgcctcctcc 1182
 5 tgggtgtcca tcagggagcc gttcgccag tgtctgtctg tctgtctgcc tctctgtctg 1242
 agggcac 1249

10 <210> 2
 <211> 20
 <212> PRT
 <213> Homo sapiens

15 <400> 2
 Met Ser Pro Ser Pro Thr Ala Leu Phe Cys Leu Gly Leu Cys Leu Gly
 1 5 10 15
 Arg Val Pro Ala
 20 20

<210> 3
 <211> 319
 25 <212> PRT
 <213> Homo sapiens

<400> 3
 Gln Ser Gly Pro Leu Pro Lys Pro Ser Leu Gln Ala Leu Pro Ser Ser
 30 1 5 10 15
 Leu Val Pro Leu Glu Lys Pro Val Thr Leu Arg Cys Gln Gly Pro Pro
 20 25 30
 35 Gly Val Asp Leu Tyr Arg Leu Glu Lys Leu Ser Ser Ser Arg Tyr Gln
 35 40 45
 Asp Gln Ala Val Leu Phe Ile Pro Ala Met Lys Arg Ser Leu Ala Gly
 50 55 60
 40 Arg Tyr Arg Cys Ser Tyr Gln Asn Gly Ser Leu Trp Ser Leu Pro Ser
 65 70 75 80
 Asp Gln Leu Glu Leu Val Ala Thr Gly Val Phe Ala Lys Pro Ser Leu
 45 85 90 95
 Ser Ala Gln Pro Gly Pro Ala Val Ser Ser Gly Gly Asp Val Thr Leu
 100 105 110
 50 Gln Cys Gln Thr Arg Tyr Gly Phe Asp Gln Phe Ala Leu Tyr Lys Glu
 115 120 125
 Gly Asp Pro Ala Pro Tyr Lys Asn Pro Glu Arg Trp Tyr Arg Ala Ser
 130 135 140
 55 Phe Pro Ile Ile Thr Val Thr Ala Ala His Ser Gly Thr Tyr Arg Cys
 145 150 155 160
 Tyr Ser Phe Ser Ser Arg Asp Pro Tyr Leu Trp Ser Ala Pro Ser Asp
 60 165 170 175

Pro Leu Glu Leu Val Val Thr Gly Thr Ser Val Thr Pro Ser Arg Leu
 180 185 190
 5 Pro Thr Glu Pro Pro Ser Ser Val Ala Glu Phe Ser Glu Ala Thr Ala
 195 200 205
 Glu Leu Thr Val Ser Phe Thr Asn Lys Val Phe Thr Thr Glu Thr Ser
 210 215 220
 10 Arg Ser Ile Thr Thr Ser Pro Lys Glu Ser Asp Ser Pro Ala Gly Pro
 225 230 235 240
 Ala Arg Gln Tyr Tyr Thr Lys Gly Asn Leu Val Arg Ile Cys Leu Gly
 245 250 255
 15 Ala Val Ile Leu Ile Ile Leu Ala Gly Phe Leu Ala Glu Asp Trp His
 260 265 270
 20 Ser Arg Arg Lys Arg Leu Arg His Arg Gly Arg Ala Val Gln Arg Pro
 275 280 285
 Leu Pro Pro Leu Pro Pro Leu Pro Gln Thr Arg Lys Ser His Gly Gly
 290 295 300
 25 Gln Asp Gly Gly Arg Gln Asp Val His Ser Arg Gly Leu Cys Ser
 305 310 315

<160> 3

30

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/03683

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/705 A61K38/17 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, STRAND, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GIBBINS JONATHAN M ET AL: "Glycoprotein VI is the collagen receptor in platelets which underlies tyrosine phosphorylation of the Fc receptor gamma-chain." FEBS LETTERS, vol. 413, no. 2, 1997, pages 255-259, XP002143941 ISSN: 0014-5793 the whole document	1
A	----- -/--	2-9



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"8" document member of the same patent family

Date of the actual completion of the international search

1 August 2000

Date of mailing of the international search report

21/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gurdjian, D

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/03683

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JANDROT-PERRUS MARTINE ET AL: "Adhesion and activation of human platelets induced by convulxin involve glycoprotein VI and integrin alpha-2-beta-1." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 272, no. 43, 1997, pages 27035-27041, XP002143942 ISSN: 0021-9258	1
A	the whole document	2-9
X	GIBBINS JONATHAN M ET AL: "The p85 subunit of phosphatidylinositol 3-kinase associated with the Fc receptor gamma-chain and linker for activator of T cells (LAT) in platelets stimulated by collagen and convulxin." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 273, no. 51, 18 December 1998 (1998-12-18), pages 34437-34443, XP002143943 ISSN: 0021-9258	1
A	the whole document	2-9
L	WO 95 11259 A (THROMBOSIS RES INST ;JEHANLI AHMED MOHAMMED TAKI (GB); PATEL GEETA) 27 April 1995 (1995-04-27) abstract; claims 1-25	1
P,X	CLEMETSON JEANNINE M ET AL: "The platelet collagen receptor glycoprotein VI is a member of the immunoglobulin superfamily closely related to FcalphaR and the natural killer receptors." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 274, no. 41, 8 October 1999 (1999-10-08), pages 29019-29024, XP002143944 ISSN: 0021-9258	1-5
P,X	the whole document	
P,X	MIURA Y: "Platelet glycoprotein VI" EMBL DATABASE ; ACCESSION NUMBER AB035073, 14 January 2000 (2000-01-14), XP002143945 the whole document	1-5

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 00/03683

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9511259 A	27-04-1995	AU 701576 B	04-02-1999
		AU 7946594 A	08-05-1995
		EP 0724599 A	07-08-1996
		JP 9506863 T	08-07-1997
		NZ 275492 A	27-05-1998
		US 6001802 A	14-12-1999
		US 5952304 A	14-09-1999
<hr/>			